arXiv:1702.01069v2 [math.MG] 16 Feb 2017

Random polytopes: variances and central limit theorems for intrinsic volumes

Christoph Thäle*, Nicola Turchi†‡ and Florian Wespi§

Abstract

Short and transparent proofs of upper variance bounds and central limit theorems for intrinsic volumes of random polytopes in smooth convex bodies are presented. They combine different tools such as estimates for floating bodies with Stein's method from probability theory.

Keywords. Central limit theorem, intrinsic volume, random polytope, stochastic geometry, variance
MSC (2010). 52A22, 60D05, 60F05

1 Introduction and main results

Fix a space dimension $n \geq 2$, let $N \geq n+1$ and suppose that $X_1, \ldots, X_N$ are independent random points that are uniformly distributed in a prescribed convex body K, which we assume to have a boundary which is twice differentiable and has positive Gaussian curvature everywhere. The convex hull of $X_1, \ldots, X_N$ is denoted by K_N. In this note we are interested in the intrinsic volumes $V_j(K_N)$ of K_N, $j \in \{1, \ldots, n\}$. These functionals are of particular importance in convex and integral geometry since they (together with the Euler-characteristic) form a basis of the vector space of all motion invariant and continuous valuations on convex bodies according to Hadwiger's celebrated theorem, cf. [13]. The purpose of this text is to provide an upper bound for the variance of $V_j(K_N)$ and to prove a central limit theorem for $V_j(K_N)$, as $N \to \infty$. Both results are not totally new. The upper variance bound for $V_j(K_N)$ has been found earlier by Bárány, Fodor and Vigh [2] by a long and very technical proof, and the central limit theorem has first been considered by Reitzner [11] in the case that $j = n$. The central limit theorem for general $j \in \{1, \ldots, n\}$ was known for a long time only in the special case that K is the n-dimensional Euclidean unit ball, see the paper of Calka, Schreiber and Yukich [5] and also that of Schreiber [14] for the situation $j = 1$. Only very recently (in parallel and independently of us) Lachièze-Rey, Schulte and Yukich [8] gave a proof for the general case by embedding the problem into the theory of so-called stabilizing functionals.

Our first contribution is a very quick and transparent proof of the upper variance bound by using estimates for floating bodies. This will lead to the right order up to a logarithmic term and extends an argument of Vu [15] for the volume to all intrinsic volumes. Using the same method in combination with a general normal approximation bound obtained by Chatterjee [6] and Lachièze-Rey and Peccati

*Faculty of Mathematics, Ruhr University Bochum, Germany. Email: christoph.thaele@rub.de
†Faculty of Mathematics, Ruhr University Bochum, Germany. Email: nicola.turchi@rub.de
§Institute of Mathematical Statistics and Actuarial Science, University of Bern, Switzerland. Email: florian.wespi@stat.unibe.ch
‡Research supported by the Research Training Group RTG2131 *High-dimensional Phenomena in Probability – Fluctuations and Discontinuity*.

[7] originating in Stein's method, our second (and in fact main) contribution is a quick, transparent and direct proof of the central limit theorem for $V_j(K_N)$ and all $j \in \{1,\dots,n\}$, as $N \to \infty$. More precisely, while the traditional methods (see [9, 11, 16]) first use a conditioning argument and prove the central limit theorem for a Poissonized version of the random polytopes, before pushing this result to the original model by de-Poissonization, we give a direct proof without making the detour just described. We also avoid this way the highly technical theory of stabilizing functionals developed in [8].

To present our results formally, we shall use the notation $a_N \ll b_N$ whenever for two sequences (a_N) and (b_N), $a_N \le c b_N$ for sufficiently large $N \ge n+1$ and some constant $c \in (0,\infty)$ not depending on N (but possibly on the space dimension n and the convex body K).

Proposition 1.1. *Let $K \subset \mathbb{R}^n$ be a convex body with twice differentiable boundary and strictly positive Gaussian curvature everywhere. Then, for all $j \in \{1,\dots,n\}$, one has that*

$$\operatorname{Var} V_j(K_N) \ll N^{-\frac{n+3}{n+1}} (\log N)^{2\frac{n+2}{n+1}}.$$

Remark 1.2. The lower and upper variance bound

$$N^{-\frac{n+3}{n+1}} \ll \operatorname{Var} V_j(K_N) \ll N^{-\frac{n+3}{n+1}} \tag{1}$$

were in fact proven by Bárány, Fodor and Vigh [2]. In particular, to achieve the lower bound they extract an event that has positive probability and on which the variance of $V_j(K_N)$ is bounded from below by a quantity that is asymptotically of the desired order. The upper bound follows by using the Efron-Stein jackknife inequality and more careful estimates for floating bodies.

Next, we define the centred and normalized intrinsic volume functionals

$$W_j(K_N) := \frac{V_j(K_N) - \mathbb{E}[V_j(K_N)]}{\sqrt{\operatorname{Var} V_j(K_N)}}, \qquad j \in \{1,\dots,n\}, N \ge n+1,$$

for which we prove the following central limit theorem. As discussed earlier, this extends the results in [5, 11] to arbitrary intrinsic volumes or to more general convex bodies, respectively.

Theorem 1.3. *Let $K \subset \mathbb{R}^n$ be a convex body with twice differentiable boundary and strictly positive Gaussian curvature everywhere. Then, for all $j \in \{1,\dots,n\}$, one has that $W_j(K_N)$ converges in distribution to a standard Gaussian random variable, as $N \to \infty$.*

Remark 1.4. Our proof will also deliver a rate of convergence in the previous central limit theorem. It is measured by the so-called Wasserstein distance and will be of order $N^{-\frac{1}{2}+\frac{1}{n+1}} (\log N)^{3+\frac{2}{n+1}}$, as $N \to \infty$, see (9) below. Since rates of convergence are not in our focus, we refer to Section 5.4 of [8] for recent results in this direction.

The rest of this note is structured as follows. In Section 2 we collect some background material in order to keep our presentation reasonably self-contained. The proof of the upper variance bound is presented in Section 3, while the proof of the central limit theorem is the content of the final Section 4.

2 Background material

Convex bodies. By a convex body we understand a compact convex subset of $\mathbb{R}^n$ that has non-empty interior. We use the symbol B^n to denote the centred Euclidean unit ball in $\mathbb{R}^n$. For points $X_1,\dots,X_N \in \mathbb{R}^n$, we write $[X_1,\dots,X_N]$ to indicate their convex hull.

Intrinsic volumes. Let $K \subset \mathbb{R}^n$ be a convex body and fix $j \in \{0, \dots, n\}$. We denote by $G(n,j)$ the Grassmannian of j-dimensional linear subspaces of $\mathbb{R}^n$, which is supplied with the unique Haar probability measure ν_j, see [13]. For $L \in G(n,j)$, we write $\mathrm{vol}_j(K|L)$ for the j-dimensional Lebesgue measure of the orthogonal projection of K onto L. Finally, we let for integers $\ell \in \mathbb{N}$, $\kappa_\ell = \pi^{\ell/2}\Gamma(1+\frac{\ell}{2})^{-1}$ be the volume of the ℓ-dimensional unit ball. Then the jth intrinsic volume of K can be defined as

$$V_j(K) := \binom{n}{j} \frac{\kappa_n}{\kappa_j \kappa_{n-j}} \int_{G(n,j)} \mathrm{vol}_j(K|L)\, \nu_j(\mathrm{d}L)\,. \tag{2}$$

For example, $V_n(K)$ is the ordinary volume (Lebesgue measure), $V_{n-1}(K)$ is half of the surface area, $V_1(K)$ is a constant multiple of the mean width and $V_0(K) \equiv 1$ is the Euler-characteristic of K.

An estimate for subspaces. Fix $j \in \{1, \dots, n-1\}$, $L \in G(n,j)$ and $z \in \mathbb{S}^{n-1}$. The angle $\sphericalangle(z,L)$ between z and L is defined as the minimum angle $\min\{\sphericalangle(z,x) : x \in L\}$. We now recall from [2, Lemma 1] that

$$\nu_j(\{L \in G(n,j) : \sphericalangle(z,L) \le a\}) \ll a^{n-j} \tag{3}$$

for all sufficiently small $a > 0$, see also [4, Lemma 10].

Floating bodies. Let $K \subset \mathbb{R}^n$ be a convex body and $t > 0$ (we shall implicitly assume that t is sufficiently small). If H is a half space of $\mathbb{R}^n$ with $V_n(K \cap H) = t$, the set $K \cap H$ is called a t-cap of K. The union of all these t-caps is the so-called wet part of K and its complement is the t-floating body of K. In what follows, we shall denote the t-floating body by $K_{(t)}$.

The Efron-Stein jackknife inequality. Let $X_1, \dots, X_N, X_{N+1}$ be independent random points that are uniformly distributed in a convex body $K \subset \mathbb{R}^n$, and put $K_N := [X_1, \dots, X_N]$ as well as $K_{N+1} := [X_1, \dots, X_{N+1}]$. For $j \in \{1, \dots, n\}$, the Efron-Stein jackknife inequality says that

$$\mathrm{Var}\, V_j(K_N) \le (N+1)\, \mathbb{E}[(V_j(K_{N+1}) - V_j(K_N))^2]\,,$$

see [10, Section 3]. In other words, the variance of $V_j(K_N)$ is bounded by the second moment of the effect that the addition of a further point X_{N+1} has on the jth intrinsic volume.

A general bound for normal approximation. Let S be a Polish space and $f : \bigcup_{k=1}^N S^k \to \mathbb{R}$ be a measurable and symmetric function acting on point configurations of at most $N \in \mathbb{N}$ points in S. For $x = (x_1, \dots, x_N) \in S^N$, we write x^i for the vector x with the ith coordinate removed and $x^{i_1 i_2}$ for the vector that arises from x by removing coordinates i_1 and i_2. Next, we define the first- and second-order difference operator applied to $f(x) = f(x_1, \dots, x_N)$ by

$$D_i f(x) := f(x) - f(x^i) \qquad \text{and} \qquad D_{i_1,i_2} f(x) := f(x) - f(x^{i_1}) - f(x^{i_2}) + f(x^{i_1 i_2})\,,$$

respectively.

We denote by $X = (X_1, \dots, X_N)$ a random vector of elements of S. Let $X', \widetilde{X}$ be independent copies of X and say that a random vector $Z = (Z_1, \dots, Z_N)$ is a recombination of $\{X, X', \widetilde{X}\}$ provided that $Z_i \in \{X_i, X_i', \widetilde{X}_i\}$ for all $i \in \{1, \dots, N\}$.

To rephrase the normal approximation bound from [6] in the form that can be deduced from [7] we define

$$\begin{aligned}
\gamma_1 &:= \sup_{(Y,Y',Z,Z')} \mathbb{E}\big[\mathbf{1}\{D_{1,2}f(Y)\neq 0\}\,\mathbf{1}\{D_{1,3}f(Y')\neq 0\}\,(D_2f(Z))^2\,(D_3f(Z'))^2\big]\,,\\
\gamma_2 &:= \sup_{(Y,Z,Z')} \mathbb{E}\big[\mathbf{1}\{D_{1,2}f(Y)\neq 0\}\,(D_1f(Z))^2\,(D_2f(Z'))^2\big]\,,\\
\gamma_3 &:= \mathbb{E}|D_1f(X)|^4\,,\\
\gamma_4 &:= \mathbb{E}|D_1f(X)|^3\,,
\end{aligned}$$

where the suprema in the definitions of γ_1 and γ_2 run over all quadruples or triples of vectors (Y,Y',Z,Z') or (Y,Z,Z') that are recombinations of $\{X,X',\widetilde{X}\}$, respectively. Next, we define $W := f(X_1,\dots,X_N)$ and assume that $\mathbb{E}W = 0$ and $\mathbb{E}W^2 < \infty$. Then, if G denotes a standard Gaussian random variable,

$$\begin{aligned}
d\bigg(\frac{W}{\sqrt{\mathrm{Var}\,W}},G\bigg) &:= \sup_h \bigg|\,\mathbb{E}\bigg[h\bigg(\frac{W}{\sqrt{\mathrm{Var}\,W}}\bigg)\bigg] - \mathbb{E}[h(G)]\,\bigg|\\
&\ll \frac{\sqrt{N}}{\mathrm{Var}\;W}\Big(\sqrt{N^2\gamma_1}+\sqrt{N\gamma_2}+\sqrt{\gamma_3}\Big) + \frac{N}{(\mathrm{Var}\;W)^{\frac{3}{2}}}\,\gamma_4\,,
\end{aligned} \tag{4}$$

where the supremum is running over all Lipschitz functions $h:\mathbb{R}\to\mathbb{R}$ with Lipschitz constant less or equal than one (this defines what is called the Wasserstein distance). In particular, if $(W_N)_{N\in\mathbb{N}}$ is a sequence of centred random variables with finite second moment such that $d(W_N/\sqrt{\mathrm{Var}\,W_N},G)\to 0$, as $N\to\infty$, then W_N converges in distribution to G.

3 Proof of Proposition 1.1

We start by re-phrasing a result of Bárány and Dalla [3], which has been proved also by Vu [15] using different techniques, see also Lemma 2.2 in [12]. Recall that K_N is the random polytope generated by N independent random points that are uniformly distributed in a convex body $K\subset\mathbb{R}^n$ having sufficiently smooth boundary (in fact, smoothness of the boundary is not needed in the next proposition).

Proposition 3.1. *For any $\beta\in(0,\infty)$, there exists a constant $c=c(\beta,n)\in(0,\infty)$ only depending on β and on n such that the event that K_N does not contain the $\frac{c\log N}{N}$-floating body is $N^{-\beta}$, whenever N is sufficiently large.*

By combining the Efron-Stein jackknife inequality with Proposition 3.1, we can now present our proof of Proposition 1.1.

Proof of Proposition 1.1. In a first step we prove the upper variance bound for $K=B^n$, the centred Euclidean ball with radius one. According to Proposition 3.1, we see that for any $\beta\in(0,\infty)$ there exists a constant $c=c(\beta,n)\in(0,\infty)$ such that $K_{(c\log N/N)}\subseteq K_N$ with probability greater or equal than $1-N^{-\beta}$. In what follows, we shall abbreviate the latter event by B. Note that we choose β large enough ($\beta=5$ will be sufficient for all our purposes) and we denote by $c=c(\beta,n)$ the constant implied by Proposition 3.1.

Now, we apply the Efron-Stein jackknife inequality, which says that

$$\operatorname{Var} V_j(K_N) \leq (N+1)\,\mathbb{E}[(V_j(K_{N+1}) - V_j(K_N))^2]\,,$$

where the random polytope K_{N+1} arises from K_N by addition of a further independent random point X_{N+1}. By conditioning on the event B introduced above (and on its complement B^c), we split the last expectation into the two terms

$$\mathbb{E}[(V_j(K_{N+1}) - V_j(K_N))^2\,\mathbf{1}_B] + \mathbb{E}[(V_j(K_{N+1}) - V_j(K_N))^2\,\mathbf{1}_{B^c}]\,.$$

On B^c we estimate the difference $(V_j(K_{N+1}) - V_j(K_N))^2$ trivially by $V_j(K)^2$, which is a constant independent of N. Since $\mathbb{P}(B^c) \leq N^{-\beta}$ and since we have chosen β sufficiently large, we obtain (regarding the exponents at N appearing in the computations below)

$$\operatorname{Var} V_j(K_N) \ll N\,\mathbb{E}[(V_j(K_{N+1}) - V_j(K_N))^2\,\mathbf{1}_B]\,.$$

Using the definition (2) of the intrinsic volumes, it follows that

$$\operatorname{Var} V_j(K_N) \ll N\,\mathbb{E}\left[\left(\int_{G(n,j)} \operatorname{vol}_j((K_{N+1}|L) \setminus (K_N|L))\,\nu_j(\mathrm{d}L)\right)^2 \mathbf{1}_B\right]. \tag{5}$$

On the event B we first notice that $\operatorname{vol}_j((K_{N+1}|L) \setminus (K_N|L))$ is zero if $X_{N+1} \in K_N$. So, we can restrict to the situation that $X_{N+1} \in K \setminus K_N$, which conditioned on B occurs with probability $V_n(K \setminus K_N) \ll V_n(K \setminus K_{(c\log N/N)}) \ll (\log N/N)^{\frac{2}{n+1}}$, cf. [1, Theorem 6.3].

It is our aim to define a full-dimensional cap C such that $K_{N+1} \setminus K_N$ is contained in C. For this reason, we define z to be the closest point to X_{N+1} on ∂K (we notice that z is uniquely determined if $K_{(c\log N/N)}$ is non-empty). The visible region of z is defined as

$$\operatorname{Vis}_z(N) := \{x \in K \setminus K_{(c\log N/N)} : [x,z] \cap K_{(c\log N/N)} = \emptyset\}\,.$$

By definition of B and the floating body $K_{(c\log N/N)}$, the diameter of $\operatorname{Vis}_z(N)$ is $c_1(\log N/N)^{\frac{1}{n+1}}$, where $c_1 \in (0,\infty)$. Let us denote by $D(z, c_1(\log N/N)^{\frac{1}{n+1}})$ the set of all points on the boundary of K which are of distance at most $c_1(\log N/N)^{\frac{1}{n+1}}$ to z. Then, it follows from [15, Lemma 6.2] that $C := \operatorname{conv}\{D(z, c_1(\log N/N)^{\frac{1}{n+1}})\}$ has volume of order at most $\log N/N$. Moreover, C is in fact a spherical cap and the central angle of it is denoted by α. For a subspace $L \in G(n,j)$, one has that $(K_{N+1}|L) \setminus (K_N|L) \subseteq C|L$. The volume $\operatorname{vol}_j(C|L)$ of the projected cap $C|L$ is $\operatorname{vol}_j(C|L) \ll (\log N/N)^{\frac{j+1}{n+1}}$. Indeed, the height of $C|L$ keeps the order of the height of C, namely $(\log N/N)^{\frac{2}{n+1}}$, while the order of its base changes from $((\log N/N)^{\frac{1}{n+1}})^{n-1}$ to $((\log N/N)^{\frac{1}{n+1}})^{j-1}$, since L is a subspace of dimension j. Note that, by construction of C, if $\sphericalangle(z,L)$, the angle between z and L, is too wide compared to α, then $C|L \subseteq K_N|L$, for sufficiently large N. In particular $(K_{N+1} \setminus K_N)|L \subseteq K_N|L$, which implies $K_{N+1}|L = K_N|L$. In fact, it is easily checked that the integrand in (5) can only be non-zero if $\sphericalangle(z,L) \ll \alpha$ (the constant can be taken to be 2 in the case of the ball). Therefore, we can restrict the integration in (5) to the set $\{L \in G(n,j) : \sphericalangle(z,L) \ll \alpha\}$. It is not difficult to verify that $\alpha \ll V_n(C)^{\frac{1}{n+1}}$, see also Equation (27) in [2].

Taken all together, this yields

$$\operatorname{Var} V_j(K_N) \ll N \cdot \left(\frac{\log N}{N}\right)^{\frac{2}{n+1}} \cdot \left(\frac{\log N}{N}\right)^{2\frac{j+1}{n+1}} \nu_j\Big(\Big\{L \in G(n,j) : \sphericalangle(z,L) \ll V_n(C)^{\frac{1}{n+1}}\Big\}\Big)^2.$$

According to (3) and the fact that $V_n(C) \ll \log N/N$, it holds that

$$\nu_j\Big(\Big\{L \in G(n,j) : \sphericalangle(z,L) \ll V_n(C)^{\frac{1}{n+1}}\Big\}\Big) \ll \left(\frac{\log N}{N}\right)^{\frac{n-j}{n+1}},$$

which in turn implies

$$\operatorname{Var} V_j(K_N) \ll N \cdot \left(\frac{\log N}{N}\right)^{\frac{2}{n+1}} \cdot \left(\frac{\log N}{N}\right)^{2\frac{j+1}{n+1}} \cdot \left(\frac{\log N}{N}\right)^{2\frac{n-j}{n+1}} = N^{-\frac{n+3}{n+1}} (\log N)^{2\frac{n+2}{n+1}}.$$

To extend the proof to the general case, we argue as in [2, Section 6]. Namely, since K is compact, we can choose $\gamma \in (0,\infty)$ and $\Gamma \in (0,\infty)$ to be, respectively, the global upper and the global lower bound on the principal curvatures of ∂K. Remark 5 on page 126 of [13] ensures that under these circumstances all projected images of K also have a boundary with the same features as ∂K, and we choose γ and Γ such that they also bound the principal curvatures of each j-dimensional projection of K. Since we can approximate ∂K locally with affine images of balls, the construction of C and the relations regarding its volume, its central angle and the subspaces L which ensure $C|L \subseteq K_N|L$ are not afflicted. Due to this, the relations $\operatorname{vol}_j(C|L) \ll (\log N/N)^{\frac{j+1}{n+1}}$, $\alpha \ll V_n(C)^{\frac{1}{n+1}} \ll (\log N/N)^{\frac{1}{n+1}}$ and $\sphericalangle(z,L) \ll \alpha$ from above still hold, but this time the implicit constants depend on γ and Γ. From here, the proof can be completed in the same way as for the ball. □

4 Proof of Theorem 1.3

In order to prepare for the proof of Theorem 1.3 let us introduce the two events B_1 and B_2 as follows. We denote by B_1 the event that the random polytope $[X_2,\dots,X_N]$ contains the floating body $K_{(c\log N/N)}$, where $c \in (0,\infty)$ is the constant from the proof of Proposition 1.1 presented in the previous section. By the discussion there, the probability $\mathbb{P}(B_1^c)$ is very small. More precisely, for sufficiently large N,

$$\mathbb{P}(B_1^c) \le (N-1)^{-\beta} \le c_1 N^{-\beta}, \tag{6}$$

where $c_1 \in (0,\infty)$ is a constant independent of N. Next, we let Y, Y', Z, Z' be recombinations of our random vector $X = (X_1,\dots,X_N)$ and denote by B_2 the event that $\bigcap_{W\in\{Y,Y',Z,Z'\}}[W_4,\dots,W_N]$ contains $K_{(c\log N/N)}$. By the union bound it follows that the probability of B_2^c is also small:

$$\mathbb{P}(B_2^c) \le c_2 N^{-\beta}, \tag{7}$$

where $c_2 \in (0,\infty)$ is again a constant independent of N.

Proof of Theorem 1.3. The idea of the proof is to apply the normal approximation bound (4) to the random variables

$$W = f(X_1,\dots,X_N) := V_j([X_1,\dots,X_N]) - \mathbb{E}V_j(K_N),$$

which clearly satisfy the assumptions made in Section 2. To this end, we need to control, in particular, the first- and second-order difference operators $D_iW = D_iV_j(K_N)$ and $D_{i_1,i_2}W = D_{i_1,i_2}V_j(K_N)$ for $i, i_1, i_2 \in \{1,\dots,N\}$.

Conditioned on the event B_1, we use (2) to estimate the first-order difference operator applied to the intrinsic volume functional $V_j(K_N)$ as follows:

$$\begin{aligned} D_1V_j(K_N) &= \binom{n}{j}\frac{\kappa_n}{\kappa_j\kappa_{n-j}}\int_{G(n,j)}\mathrm{vol}_j((K_N|L)\setminus([X_2,\dots,X_N]|L))\,\nu_j(\mathrm{d}L) \\ &\qquad\qquad \times \mathbf{1}\{X_1\in K\setminus K_{(c\log N/N)}\} \\ &\ll \left(\frac{\log N}{N}\right)^{\frac{j+1}{n+1}}\cdot\left(\frac{\log N}{N}\right)^{\frac{n-j}{n+1}}\mathbf{1}\{X_1\in K\setminus K_{(c\log N/N)}\} \\ &\ll \frac{\log N}{N}\,\mathbf{1}\{X_1\in K\setminus K_{(c\log N/N)}\}\,. \end{aligned} \tag{8}$$

Note that we used here the same arguments as in the proof of Proposition 1.1. Moreover, on the complement B_1^c of B_1, we simply use the trivial estimate $D_1V_j(K_N)\le V_j(K)$ and thus conclude that

$$\begin{aligned} \mathbb{E}[(D_1V_j(K_N))^p] &= \mathbb{E}[(D_1V_j(K_N))^p\,\mathbf{1}_{B_1}]+\mathbb{E}[(D_1V_j(K_N))^p\,\mathbf{1}_{B_1^c}] \\ &\ll \left(\frac{\log N}{N}\right)^p V_n(K\setminus K_{(c\log N/N)}) \ll \left(\frac{\log N}{N}\right)^{p+\frac{2}{n+1}} \end{aligned}$$

for all $p\in\{1,2,3,4\}$, where we applied the probability estimate (6) in the second step, which ensures that the second term can be made very small for large N (the choice for p is motived by our applications below). As a consequence, we can already bound the terms appearing in the normal approximation bound (4) that involve γ_3 and γ_4. Namely, using the lower variance bound (1) we see that

$$\begin{aligned} \frac{\sqrt{N}}{\mathrm{Var}\,V_j(K_N)}\sqrt{\gamma_3} &\ll \frac{\sqrt{N}}{N^{-\frac{n+3}{n+1}}}\left(\frac{\log N}{N}\right)^{2+\frac{1}{n+1}} = N^{-\frac12+\frac{1}{n+1}}(\log N)^{2+\frac{1}{n+1}}\,, \\ \frac{N}{(\mathrm{Var}\,V_j(K_N))^{\frac32}}\gamma_4 &\ll \frac{N}{N^{-\frac32\frac{n+3}{n+1}}}\left(\frac{\log N}{N}\right)^{3+\frac{2}{n+1}} = N^{-\frac12+\frac{1}{n+1}}(\log N)^{3+\frac{2}{n+1}}\,. \end{aligned}$$

Next, we consider the second-order difference operator. We define, for $z\in K\setminus K_{(c\log N/N)}$,

$$\mathrm{Vis}_z(N) := \{x\in K\setminus K_{(c\log N/N)} : [x,z]\cap K_{(c\log N/N)}=\emptyset\}\,.$$

On the event B_2 it may be concluded from (8) that $D_if(V)^2\ll(\log N/N)^2$ for all $i\in\{1,2,3\}$ and $V\in\{Z,Z'\}$. Next, we note that on B_2 the following inclusion holds:

$$\begin{aligned} \{D_{1,2}f(Y)\neq0\} &\subseteq \{Y_1\in K\setminus K_{(c\log N/N)}\}\cap\{Y_2\in K\setminus K_{(c\log N/N)}\}\cap\{\mathrm{Vis}_{Y_1}(N)\cap\mathrm{Vis}_{Y_2}(N)\neq\emptyset\} \\ &\subseteq \{Y_1\in K\setminus K_{(c\log N/N)}\}\cap\Big\{Y_2\in\bigcup_{x\in\mathrm{Vis}_{Y_1}(N)}\mathrm{Vis}_x(N)\Big\}\,. \end{aligned}$$

The same applies to $D_{1,3}f(Y')$ as well. We thus infer that

$$\begin{aligned} &\mathbb{E}\big[\mathbf{1}\{D_{1,2}f(Y)\neq0\}\,\mathbf{1}_{B_2}\big] \\ &\le \mathbb{P}\big(Y_1\in K\setminus K_{(c\log N/N)}\big)\,\mathbb{P}\Big(Y_2\in\bigcup_{x\in\mathrm{Vis}_{Y_1}(N)}\mathrm{Vis}_x(N)\;\Big|\;Y_1\in K\setminus K_{(c\log N/N)}\Big) \end{aligned}$$

$$\leq \mathbb{P}\big(Y_1 \in K \setminus K_{(c\log N/N)}\big) \sup_{z\in K\setminus K_{(c\log N/N)}} \mathbb{P}\Big(Y_2 \in \bigcup_{x\in \mathrm{Vis}_z(N)} \mathrm{Vis}_x(N)\Big)$$
$$= V_n\big(K \setminus K_{(c\log N/N)}\big) \sup_{z\in K\setminus K_{(c\log N/N)}} V_n\Big(\bigcup_{x\in \mathrm{Vis}_z(N)} \mathrm{Vis}_x(N)\Big).$$

Since the diameter of the previous union is of order $(\log N/N)^{\frac{1}{n+1}}$, it follows from [15, Lemma 6.2] that

$$\Delta(N) := \sup_{z\in K\setminus K_{(c\log N/N)}} V_n\Big(\bigcup_{x\in \mathrm{Vis}_z(N)} \mathrm{Vis}_x(N)\Big) \ll \frac{\log N}{N}.$$

Moreover, on the complement B_2^c of B_2 we estimate all the indicator functions by one and the value of all difference operators by the constant $V_j(K)$. Since $\mathbb{P}(B_2^c)$ is small in N, this readily implies

$$\gamma_2 \ll \left(\frac{\log N}{N}\right)^4 V_n(K \setminus K_{(c\log N/N)})\,\Delta(N) \ll \left(\frac{\log N}{N}\right)^{5+\frac{2}{n+1}}.$$

Analogously, we can bound γ_1. First, suppose that $Y_1 = Y_1'$. Then, conditioned on B_2,

$$\begin{aligned}
&\{D_{1,2}f(Y) \neq 0\} \cap \{D_{1,3}f(Y') \neq 0\}\\
&\subseteq \{\{Y_1, Y_2, Y_3'\} \subseteq K \setminus K_{(c\log N/N)}\} \cap \{\mathrm{Vis}_{Y_2}(N) \cap \mathrm{Vis}_{Y_1}(N) \neq \emptyset\} \cap \{\mathrm{Vis}_{Y_3'}(N) \cap \mathrm{Vis}_{Y_1}(N) \neq \emptyset\}\\
&\subseteq \{Y_1 \in K \setminus K_{(c\log N/N)}\} \cap \Big\{\{Y_2, Y_3'\} \subseteq \bigcup_{x\in \mathrm{Vis}_{Y_1}(N)} \mathrm{Vis}_x(N)\Big\},
\end{aligned}$$

and arguing as before leads to

$$\begin{aligned}
&\mathbb{E}\big[\mathbf{1}\{D_{1,2}f(Y) \neq 0\}\,\mathbf{1}\{D_{1,3}f(Y') \neq 0\}\,\mathbf{1}_{B_2}\big]\\
&\leq \mathbb{P}\big(Y_1 \in K \setminus K_{(c\log N/N)}\big) \sup_{z\in K\setminus K_{(c\log N/N)}} \mathbb{P}\Big(\{Y_2, Y_3'\} \subseteq \bigcup_{x\in \mathrm{Vis}_z(N)} \mathrm{Vis}_x(N)\Big)\\
&\leq V_n(K \setminus K_{(c\log N/N)})\,\Delta(N)^2.
\end{aligned}$$

Note that the case $Y_1 \neq Y_1'$ gives a smaller order since, by independence, it leads to an extra factor $V_n(K \setminus K_{(c\log N/N)})$. Thus, by conditioning on B_2 and its complement, we obtain

$$\gamma_1 \ll \left(\frac{\log N}{N}\right)^4 V_n(K \setminus K_{(c\log N/N)})\,\Delta(N)^2 \ll \left(\frac{\log N}{N}\right)^{6+\frac{2}{n+1}}.$$

Now, the other terms appearing in the normal approximation bound (4) can be estimated by

$$\frac{\sqrt{N}}{\mathrm{Var}\, V_j(K_N)}\sqrt{N^2\gamma_1} \ll \frac{\sqrt{N}}{N^{-\frac{n+3}{n+1}}}\sqrt{N^2\cdot\left(\frac{\log N}{N}\right)^{6+\frac{2}{n+1}}} = N^{-\frac{1}{2}+\frac{1}{n+1}}(\log N)^{3+\frac{1}{n+1}},$$
$$\frac{\sqrt{N}}{\mathrm{Var}\, V_j(K_N)}\sqrt{N\gamma_2} \ll \frac{\sqrt{N}}{N^{-\frac{n+3}{n+1}}}\sqrt{N\cdot\left(\frac{\log N}{N}\right)^{5+\frac{2}{n+1}}} = N^{-\frac{1}{2}+\frac{1}{n+1}}(\log N)^{\frac{5}{2}+\frac{1}{n+1}}.$$

Putting together all estimates, we arrive at

$$d\Big(\frac{V_j(K_N)-\mathbb{E}V_j(K_N)}{\sqrt{\operatorname{Var}V_j(K_N)}},G\Big) \ll N^{-\frac{1}{2}+\frac{1}{n+1}}\Big((\log N)^{3+\frac{1}{n+1}}+(\log N)^{\frac{5}{2}+\frac{1}{n+1}} + (\log N)^{2+\frac{1}{n+1}}+(\log N)^{3+\frac{2}{n+1}}\Big) \ll N^{-\frac{1}{2}+\frac{1}{n+1}}(\log N)^{3+\frac{2}{n+1}} \tag{9}$$

from the normal approximation bound (4). In particular, as $N\to\infty$, this converges to zero and so the random variables

$$W_j(K_N)=\frac{V_j(K_N)-\mathbb{E}V_j(K_N)}{\sqrt{\operatorname{Var}V_j(K_N)}}$$

converge in distribution to the standard Gaussian random variable G. The proof of Theorem 1.3 is thus complete. □

Acknowledgement

We would like to thank Matthias Schulte for pointing us to a mistake in the first version of this manuscript.

References

[1] I. Bárány (2007): Random points and lattice points in convex bodies. Bulletin of the AMS **45**, 339–365.

[2] I. Bárány, F. Fodor and V. Vigh (2010). Intrinsic volumes of inscribed random polytopes in smooth convex bodies. Adv. Appl. Probab. **42**, 605–619.

[3] I. Bárány and L. Dalla (1997). Few points to generate a random polytope. Mathematika **44 44**, 325–331.

[4] I. Bárány and C. Thäle (2016): Intrinsic volumes and Gaussian polytopes: the missing piece of the jigsaw. arXiv: 1606.07945.

[5] P. Calka, T. Schreiber and J.E. Yukich (2013): Brownian limits, local limits and variance asymptotics for convex hulls in the ball. Ann. Probab. **41**, 50–108.

[6] S. Chatterjee (2008): A new method of normal approximation. Ann. Probab. **36**, 1584–1610.

[7] R. Lachièze-Rey and G. Peccati (2016+): New Berry-Esseen bounds for functionals of binomial point processes. to appear in Ann. Appl. Probab.

[8] R. Lachièze-Rey, M. Schulte and Y.E. Yukich (2017): Normal approximation for sums of stabilizing functionals. arXiv: 1702.00726.

[9] J. Pardon (2012): Central limit theorems for uniform model random polygons. J. Theor. Probab. **25**, 823–833.

[10] M. Reitzner (2003): Random polytopes and the Efron-Stein jackknife inequality. Ann. Probab. **31**, 2136–2166.

[11] M. Reitzner (2005): Central limit theorems for random polytopes. Probab. Theory Relat. Fields **133**, 483–507.

[12] M. Reitzner (2008): Random polytopes. In *New Perspectives in Stochastic Geometry* (edited by I. Molchanov and W.S. Kendall), 45–76, Oxford University Press.

[13] R. Schneider (2014): *Convex Bodies – The Brunn-Minkowski Theory*. 2nd expanded edition, Cambridge University Press.

[14] T. Schreiber (2002): Variance asymptotics and central limit theorems for volumes of unions of random closed sets. Adv. Appl. Probab. **34**, 520–539.

[15] V.H. Vu (2005): Sharp concentration of random polytopes. Geom. Funct. Anal. **15**, 1284–1318.

[16] V.H. Vu (2006): Central limit theorems for random polytopes in a smooth convex set. Adv. Math. **207**, 221–243.